arXiv:math/9805001v3 [math.RT] 18 Jun 1998

APPROXIMATE REPRESENTATIONS AND VIRASORO ALGEBRA

D.V.Juriev[1]

Research Center for Mathematical Physics and Informatics “Thalassa Aitheria”,
ul.Miklukho-Maklaya 20-180, Moscow 117437 Russia.
E-mail: denis@juriev.msk.ru

February, 1997 math.RT/9805001

This article being addressed, in general, to the specialists in the theory of representations of finite dimensional reductive and infinite dimensional $\mathbb{Z}$–graded Lie algebras, the theory of approximations in functional analysis and asymptotical methods in the operator calculus and, partially, in the mathematical physics (algebraic quantum field theory), is devoted to an investigation of algebraic and analytic structure of infinite dimensional hidden symmetries in the theory of representation of finite dimensional reductive Lie algebras. More precisely, the subject of the paper is a realization of the infinite dimensional $\mathbb{Z}$–graded Witt algebra of all Laurent polynomial vector fields on a circle and its one-dimensional nontrivial central extension (the Virasoro algebra) by hidden symmetries in the Verma modules over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ (the so-called q_R–conformal symmetries). These infinite dimensional Lie algebras are realized by the q_R–conformal symmetries approximately. In the article there are considered two kinds of approximate representations: the representations up to a class $\mathfrak{S}$ of operators (Hilbert-Schmidt operators, compact operators or finite-rank operators) and asymptotic representations “mod $O(\hbar)$”, where $\hbar$ is a parameter (in this case several definitions of an operator “O–large” by the parameter $\hbar$ are possible). The approximate representations of the first kind may be naturally considered in the context of the pseudo-differential calculus [1,2], whereas ones of the second kind – in the context of the asymptotic methods [3-5]. Note that the asymptotic representations “mod $O(\hbar)$” were explored in the formalism of the Karasev-Maslov asymptotic quantisation [6], and representations up to $\mathfrak{S}$ are certain generalisations of the projective representations [7,8].

Results of the article, some of which were anonced in the electronic preprints of the electronic archive on functional analysis (the Los Alamos National Laboratories, USA), should be considered more as observations than as theorems with laborious proofs, which are not clear from their formulations. Proofs of all facts are performed really in one step and are bulky computative exercises (omitted where it is possible) based on the formulas derived by the author earlier; thus, the attention should be devoted, in general, to the formulations of results themselves.

---

[1] This is an English translation of the original Russian version, which is located at the end of the article as an appendix. In the case of any differences between English and Russian versions caused by a translation the least has the priority as the original one.

§1. Preliminary definitions

1.1. The Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ and Verma modules over it. Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ is a three-dimensional space of $2{\times}2$ complex matrices with zero trace supplied by the standard commutator $[X,Y]=XY-YX$, where the right hand side multiplication is the standard matrix multiplication. In the basis

$$l_{-1}=\begin{pmatrix}0&1\\0&0\end{pmatrix},\quad l_0=\begin{pmatrix}\frac12&0\\0&-\frac12\end{pmatrix},\quad l_1=\begin{pmatrix}0&0\\-1&0\end{pmatrix}$$

the commutation relations are of the form: $[l_i,l_j]=(i-j)l_{i+j}$ $(i,j=-1,0,1)$. Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ is $\mathbb{Z}$–graded: $\deg(l_i)=-\operatorname{ad}(l_0)l_i=i$, where $\operatorname{ad}(X)$ is the adjoint action operator in the Lie algebra: $\operatorname{ad}(X)Y=[X,Y]$. Therefore, $\mathbb{Z}$–graded modules over $\mathfrak{sl}(2,\mathbb{C})$ are l_0–diagonal. A vector v in a $\mathbb{Z}$–graded module over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ is called extremal iff $l_1v=0$ and the linear span of vectors $l_{-1}^n v$ $(n\in\mathbb{Z}_+)$ coincides with the module itself (i.e. v is a cyclic vector). A $\mathbb{Z}$–graded module with an extremal vector (in this case it is defined up to a multiplier) is called extremal [11]. An extremal module is called the Verma module [12] iff the action of l_{-1} is free in it, i.e. the vectors $l_{-1}^n v$ are linearly independent. In the case of the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ the Verma modules are just the same as infinite dimensional extremal modules. An extremal weight of the Verma module is the number defined by the equality $l_0v=hv$, where v is the extremal vector. The Verma modules are defined for all complex numbers h and are pairwise nonisomorphic. Below we shall consider the Verma modules with real extremal weights only.

The Verma module V_h over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ with the extremal weight h may be realized in the space $\mathbb{C}[z]$ of polynomials of a complex variable z. The formulas for the generators of the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ are of the form:

$$l_{-1}=z,\quad l_0=z\partial_z+h,\quad l_1=z\partial_z^2+2h\partial_z,$$

here $\partial_z=\frac{d}{dz}$.

The Verma module is nondegenerate (i.e. does not contain any proper submodule) iff $h\neq-\frac{n}{2}$ $(n\in\mathbb{Z}_+)$. The Verma module V_h is called unitarizable (or hermitean) iff it admits a structure of the pre-Hilbert space such that $l_i^*=l_{-i}$. The completion of the unitarizable Verma module will be denoted by V_h^{Hilb}. The Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ acts in V_h^{Hilb} by the unbounded operators. Also it is rather useful to consider the formal Verma modules V_h^{form}, which are realized in the space $\mathbb{C}[[z]]$ of formal power series of a complex variable z, whereas the formulas for generators of the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ coincide with ones above. Note that $V_h\subseteq V_h^{\mathrm{Hilb}}\subseteq V_h^{\mathrm{form}}$ and modules V_h, V_h^{Hilb}, V_h^{form} form the Gelfand triple or the Dirac equipment of the Hilbert space V_h^{Hilb}. An action of the real form of the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ generated by the anti-Hermitean operators il_0, l_1-l_{-1} and $i(l_1+l_{-1})$ in the Hilbert space V_h^{Hilb} by the unbounded operators is exponentiated to a unitary representation of the corresponding simply connected Lie group.

In the nonunitarizable Verma module over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ there exists the unique (up to a scalar multiple) indefinite sesquilinear form $(\cdot,\cdot)$ such that $(l_iv_1,v_2)=(v_1,l_{-i}v_2)$ for any two vectors v_1 and v_2 from the Verma module. If this sesquilinear form is nondegenerate (in this case the Verma module is nondegenerate) then it has a signature (n,∞), where n is finite, and therefore, there is defined a Pontryagin completion [10] of the Verma module. The corresponding module in which the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ acts by the unbounded operators will be denoted by V_h^{Pontr}. The following chain of inclusions holds: $V_h \subseteq V_h^{\text{Pontr}} \subseteq V_h^{\text{form}}$. An action of the real form of the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ generated by the anti-Hermitean (with respect to the nondegenerate indefinite sesquilinear form $(\cdot,\cdot)$) operators il_0, $l_1 - l_{-1}$ and $i(l_1 + l_{-1})$ in the Pontryagin space V_h^{Pontr} by the unbounded operators is exponentiated to a pseudounitary representation of the corresponding simply connected Lie group.

1.2. Hidden symmetries in Verma modules over the lie algebra $\mathfrak{sl}(2,\mathbb{C})$: Lobachevskii-Berezin C^*–algebra and q_R–conformal symmetries.

Proposition 1 [13]. *In the nondegenerate Verma module V_h over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ there are uniquely defined the operators D and F such that*

$$[D,l_{-1}] = 1,\quad [D,l_0] = D,\quad [D,l_1] = D^2,$$
$$[l_{-1},F] = 1,\quad [l_0,F] = F,\quad [l_1,F] = F^2.$$

If the Verma modules are realized in the space $\mathbb{C}[z]$ of polynomials of a complex variable z then

$$D = \partial_z,\qquad F = z\tfrac{1}{\xi+2h},$$

where $\xi = z\partial_z$. The operators F and D obey the following relations:

$$[FD,DF] = 0,\qquad [D,F] = q_R(1-DF)(1-FD),$$

where $q_R = \frac{1}{2h-1}$. In the unitarizable Verma module ($q_R \neq 0$) the operators F and D are bounded and $F^* = D$, $D^* = F$.

The algebra generated by the variables t and t^* with the relations $[tt^*, t^*t] = 0$ and $[t,t^*] = q_R(1-tt^*)(1-t^*t)$ being the Berezin quantization of the Lobachevskii plane realized in the unit complex disc (the Poincarè realization) [14] is called the Lobachevskii-Berezin algebra. Proposition 1 allows to consider the Lobachevskii-Berezin algebra as a C^*–algebra. The Lobachevskii-Berezin C^*–algebra war rediscovered recently by S.Klimek and A.Lesnievsky [15].

Proposition 2 [13]. *In the nongenerate Verma module V_h over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ there are uniquely defined the operators L_n ($n \in \mathbb{Z}$) such that*

$$[l_i,L_n] = (i-n)L_{i+n}\qquad (i = 1,2,1;\ n \in \mathbb{Z}),$$

moreover, $L_i = l_i$ ($i = -1,0,1$). If the Verma modules are realized in the space $\mathbb{C}[z]$ of polynomials of a complex variable z then

$$L_k = (xi + (k+1)h)\partial_z^k\ (k \geq 0),\quad L_{-k} = z^k\tfrac{\xi+(k+1)h}{(\xi+2h)\ldots(\xi+2h+k-1)}\ (k \geq 1),$$

where $\xi = z\partial_z$. The operators L_n obey the following relations:

$$[L_n,L_m] = (n-m)L_{n+m},\ \text{if } n,m \geq -1 \text{ or } n,m \leq 1.$$

In the unitarizable Verma module the operators L_n *are unbounded and* $L_i^* = L_{-1}$.

The operators L_n are called the q_R–conformal symmetries. They may be *symbolically* represented in the form:

$$L_n = D^{nh} L_0 D^{n(1-h)}, \qquad L_{-n} = F^{n(1-h)} L_0 F^{nh}.$$

To supply the symbolical recording by a sense one should use the general commutation relations

$$[L_n, f(D)] = (-D)^{n+1} f'(D) \ (n \geq -1), \qquad [L_{-n}, f(F)] = F^{n+1} f'(D) \ (n \geq -1)$$

for $n = 0$.

The commutation relations for the operators D, F and the generators of q_R–conformal symmetries with the generators of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ mean that the families J_k and L_k $(k \in \mathbb{Z})$, where $J_i = D^i$, $J_{-i} = F^i$ $(i \in \mathbb{Z}_+)$, are families of tensor operators [8,16] for the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$.

To the families of tensor operators J_k and L_k one may correspond the generating functions – the operator fields, i.e. the formal Laurent series of a complex variable u with operator coefficients:

$$J(u) = \sum_{i \in \mathbb{Z}} J_i (-u)^{-1-i}, \quad T(u) = \sum_{i \in \mathbb{Z}} L_i (-u)^{-2-i}.$$

For any value of u an operator field realizes a mapping from V_h into V_h^{form}. The fact that J_k and T_k $(k \in \mathbb{Z})$ form families of tensor operators on the language of operator fields means that

$$[l_i, J(u)] = (-u)^i J(u) - (-u)^{i+1} J'(u), \quad [l_i, T(u)] = 2(-u)^i T(u) - (-u)^{i+1} T'(u).$$

The operator fields $V(u)$, which transform as s–differentials under the action of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ (i.e. which obey the identity $[l_i, V(u)] = s(-u)^i V(u) - (-u)^{i+1} V'(u)$), are called the $\mathfrak{sl}(2, \mathbb{C})$–primary fields of spin s; so the operator fields $J(u)$ and $T(u)$ are $\mathfrak{sl}(2, \mathbb{C})$–primary ones of spin 1 and 2. The operator fields $J(u)$ and $T(u)$ as well as their properties are explored in detail in [13].

1.3. Infinite dimensional $\mathbb{Z}$–graded Lie algebras: the Witt algebra $\mathfrak{w}^{\mathbb{C}}$ of all Laurent polynomial vector fields on a circle and the Virasoro algebra $\mathfrak{vir}^{\mathbb{C}}$, its one-dimensional nontrivial central extension. The Lie algebra $\mathrm{Vect}(S^1)$ is realized in the space of C^∞–smooth vector fields $v(t)\partial_t$ on a circle $S^1 \simeq \mathbb{R}/2\pi\mathbb{Z}$ with the commutator

$$[v_1(t)\partial_t, v_2(t)\partial_t] = (v_1(t)v_2'(t) - v_1'(t)v_2(t))\partial_t.$$

In the basis $s_n = \sin(nt)\partial_t$, $c_n = \cos(nt)\partial_t$, $h = \partial_t$ the commutation relations have the form:

$$\begin{aligned}
[s_n, s_m] &= \tfrac{1}{2}((m-n)s_{n+m} + \operatorname{sgn}(n-m)(n+m)s_{|n-m|}), \\
[c_n, c_m] &= \tfrac{1}{2}((n-m)s_{n+m} + \operatorname{sgn}(n-m)(n+m)s_{|n-m|}), \\
[s_n, c_m] &= \tfrac{1}{2}((m-n)c_{n+m} - (m+n)c_{|n-m|}) - n\delta_{nm}h, \\
[h, s_n] &= nc_n, \quad [h, c_n] = -ns_n.
\end{aligned}$$

Let us denote by $\mathrm{Vect}^{\mathbb{C}}(S^1)$ the complexification of the Lie algebra $\mathrm{Vect}(S^1)$. In the basis $e_n = ie^{ikt}\partial_t$ the commutation relations in the Lie algebra Li $\mathrm{Vect}^{\mathbb{C}}(S^1)$ have the form:

$$[e_j, e_k] = (j-k)e_{j+k}.$$

It is rather convenient to consider an imbedding of the circle S^1 into the complex plane $\mathbb{C}$ with the coordinate z, so that $z = e^{it}$ on the circle and the elements of the basis e_k ($k \in \mathbb{Z}$) are represented by the Laurent polynomial vector fields: $e_k = z^{k+1}\partial_z$. The $\mathbb{Z}$–graded Lie algebra generated by the Laurent polynomial vector fields (i.e. by the finite linear combinations of elements of the basis e_k) is called the Witt algebra and is denoted by $\mathfrak{w}^{\mathbb{C}}$. The Witt algebra $\mathfrak{w}^{\mathbb{C}}$ is the complexification of the subalgebra $\mathfrak{w}$ of the algebra $\mathrm{Vect}(S^1)$ generated by the trigonometric polynomial vector fields on a circle S^1, i.e. by the finite linear combinations of elements of the basis s_n, c_n and h.

The Lie algebra $\mathrm{Vect}(S^1)$ admits a nontrivial one-dimensional central extension defined by the Gelfand-Fuchs 2-cocycle [17]:

$$c(v_1(t)\partial_t, v_2(t)\partial_t) = \int_0^{2\pi} (v_1'(t)v_2''(t) - v_2'(t)v_1'(t))\,dt.$$

This extension being continued to the complexification $\mathrm{Vect}^{\mathbb{C}}(S^1)$ of the Lie algebra $\mathrm{Vect}(S^1)$ and reduced to the subalgebra $\mathfrak{w}^{\mathbb{C}}$ defines a central one-dimensional extension of the Witt algebra, which is called the Virasoro algebra and is denoted by $\mathfrak{vir}^{\mathbb{C}}$. The Virasoro algebra is generated by the elements e_k ($k \in \mathbb{Z}$) and the central element c with the commutation relations:

$$[e_j, e_k] = (j-k)e_{j+k} + \frac{j^3 - j}{12}c$$

and is the complexification of a central extension $\mathfrak{vir}$ of the Lie algebra $\mathfrak{w}$. In the irreducible representation the central element c of the Virasoro algebra is mapped to a scalar operator, which is proportional to the identity operator with a coefficient called *the central charge.*

§2. $\mathcal{HS}$–projective representation of the Witt algebra in Verma modules over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$

2.1. $\mathfrak{A}$–projective representations [9].

Definition 1A. Let $\mathfrak{A}$ be an associative algebra represented by linear operators in the linear space H and $\mathfrak{g}$ be the Lie algebra. The linear mapping $T : \mathfrak{g} \mapsto \mathrm{End}(H)$ is called *a $\mathfrak{A}$–projective representation* iff for any X and Y from $\mathfrak{g}$ there exists an element of the algebra $\mathfrak{A}$ represented by the operator A_{XY} such that

$$[T(X), T(Y)] - T([X,Y]) = A_{XY}.$$

If H is infinite dimensional then the representation may be realized by the unbounded operators with suitable domains of definition.

Remark 1. Definition 1A may be generalized for arbitrary anticommutative algebras. In this situation it is deeply related to the constructions of representations of the anti-commutative algebras $\mathfrak{jl}(2,\mathbb{C})$ and $\mathfrak{sl}^*(2,\mathbb{C})$ from [18,19;20:§2]. In general, it should be considered in the context of old ideas of A.I.Maltsev on representations of arbitarary nonassociative algebras [21]. General anti-commutative algebras and constructions of their $\mathfrak{A}$–projective representations are interesting for the theory of quasi-Hopf algebras, which are nonassociative as coalgebras, jacobian and c-jacobian quasi-bialgebras and related structures (see refs in [9]).

Example. If $(\mathfrak{g},\mathfrak{h})$ is a reductive pair then any representation of the Lie algebra $\mathfrak{g}$ is a $\mathcal{U}(\mathfrak{h})$–projective representation of the binary anticommutative algebra $\mathfrak{p}$ ($\mathfrak{g}=\mathfrak{h}\oplus\mathfrak{p}$, the binary operation in $\mathfrak{p}$ has the standard form: $[X,Y]_{\mathfrak{p}}=\pi([X,Y])$, where $[\cdot,\cdot]$ is the commutator in $\mathfrak{g}$ and π is the projector of $\mathfrak{g}$ onto $\mathfrak{p}$ along $\mathfrak{h}$ [22]).

Remark 2. The standard projective representation is a particular case of Def.1A if the algebra $\mathfrak{A}$ is one-dimensional and acts in H by scalar operators.

Remark 3. If H is a Hilbert (or pre-Hilbert) space then one may consider the algebra $\mathcal{HS}$ of all Hilbert-Schmidt operators as $\mathfrak{A}$. It is possible to consider the algebras $\mathcal{B}$, $\mathcal{K}$, $\mathcal{TC}$ and $\mathcal{FR}$ of all bounded, compact, trace class and finite-rank operators, too.

Definition 1B. Let $\mathfrak{A}$ be an associative algebra with an involution $*$ symmetrically represented in the Hilbert space H. If $\mathfrak{g}$ is a Lie algebra with an involution $*$ then its $\mathfrak{A}$–projective representation T in the space H is called *symmetric* iff for all elements a from $\mathfrak{g}$ $T(a^*)=T^*(a)$. Let $\mathfrak{g}$ be $\mathbb{Z}$–graded Lie algebra ($\mathfrak{g}=\oplus_{n\in\mathbb{Z}}\mathfrak{g}_n$) with an involution $*$ such that $\mathfrak{g}_n^*=\mathfrak{g}_{-n}$ and involution is identical on the subalgebra $\mathfrak{g}_0$. Let us extend the $\mathbb{Z}$–grading and the involution $*$ from the Lie algebra $\mathfrak{g}$ to the tensor algebra $\mathbf{T}^{\cdot}(\mathfrak{g})$. A symmetric $\mathfrak{A}$–projective representation of $\mathfrak{g}$ ia called *absolutely symmetric* iff for any element a of the algebra $\mathbf{T}^{\cdot}(\mathfrak{g})$ such that $\deg(a)=0$ the identity $T(a)=T^*(a)$ holds (here the representation T of the Lie algebra $\mathfrak{g}$ in H is extended to the mapping of $\mathbf{T}^{\cdot}(\mathfrak{g})$ into $\mathrm{End}(H)$.

Definition 1C. A $\mathfrak{A}$–projective representation T of the Lie algebra $\mathfrak{g}$ in the linear space H is called *almost absolutely closed* iff for any natural n and arbitrary elements $X_0,X_1,X_2,\ldots X_{n+1}$ of the Lie algebra $\mathfrak{g}$ there exists an element $\varphi(X_0,X_1,X_2,\ldots X_{n+1})$ of the algebra $\mathfrak{g}$ such that

$$[\ldots[[T(X_0),T(X_1)],T(X_2)],\ldots,T(X_{n+1})]\equiv T(\varphi(X_0,X_1,X_2,\ldots X_{n+1}))\pmod{\mathfrak{A}},$$

here $\mathfrak{A}$ is considered as represented in $\mathrm{End}(H)$. An almost absolutely closed $\mathfrak{A}$–projective representation T of the Lie algebra $\mathfrak{g}$ in the linear space H is called *absolutely closed* iff $\varphi(\cdot,\ldots,\cdot)\equiv 0$.

The mappings $(X_0,X_1,X_2,\ldots X_{n+1})\mapsto\varphi(X_0,X_1,X_2,\ldots X_{n+1})$ associated with an almost absolutely closed $\mathfrak{A}$–projective representation of the Lie algebra $\mathfrak{g}$ define the higher brackets in the Lie algebra $\mathfrak{g}$. Objects with higher brackets systematically appear in many branches of mathematics and mathematical physics (see e.g. the book [6] or the article [23] among others and numerous references wherein).

2.2. $\mathcal{HS}$–projective representation of the Witt algebra by q_R–conformal symmetries in the unitarizable Verma modules V_h over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ [9]. Note that the Witt algebra $\mathfrak{w}^{\mathbb{C}}$ admits a natural involution $*$.

Theorem 1A. *The generators L_k ($k\in\mathbb{Z}$) of q_R–conformal symmetries in the unitarizable Verma module V_h over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ realize an absolutely* *symmetric $\mathcal{HS}$–projective representation of the Witt algebra $\mathfrak{w}^{\mathbb{C}}$. Adding the tensor operators J_k ($k\in\mathbb{Z}$) to the tensor operators L_k one receives an absolutely symmetric $\mathcal{HS}$–projective representation of a semi-direct sum of the Witt algebra and the infinite-dimensional $\mathbb{Z}$–graded Heisenberg algebra (the one-dimensional central extension of the infinite-dimensional $\mathbb{Z}$–graded abelian Lie algebra $\mathbb{C}[z,z^{-1}]$ of Laurent polynomials).*

The statement of the theorem follows from the explicit formulas for generators of q_R–conformal symmetries and tensor operators J_k. A verification on the belonging to the class $\mathcal{HS}$ of Hilbert-Schmidt operators for operators of the fixed degree with respect to the grading in the $\mathbb{Z}$–graded space of polynomials $\mathbb{C}[z]$ supplied by the certain scalar product with orthogonal weight spaces (such as any unitarizable Verma module over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ is) does not produce any problems.

Remark 4. The $\mathcal{HS}$–projective representations of the Witt algebra in the unitarizable Verma modules over $\mathfrak{sl}(2,\mathbb{C})$ are absolutely closed.

Remark 5. Theorem 1A is generalized for the pseudo-unitary case with the substitution of the class $\mathcal{HS}$ of Hilbert-Schmidt operators to the class $\mathcal{K}$ of compact operators.

The results on the "exponentiated" version of Theorem 1A are announced in the e-prints [24,25].

2.3. $\mathcal{FR}$–projective representations of the Witt algebra in Verma modules over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ with the extremal weight 1 and $\frac{1}{2}$. The sattement of Theorem 1A may be strengthened in some particular cases.

Theorem 1B. *For $h=1$ or $h=\frac{1}{2}$ the $\mathcal{HS}$–projective representation of the Witt algebra by q_R–conformal symmetries in the unitarizable Verma modules over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ is $\mathcal{FR}$–projective. For $h=\frac{1}{2}$ the $\mathcal{HS}$–projective representation of a semi-direct sum of the Witt algebra and the infinite-dimensional Heisenberg algebra is also $\mathcal{FR}$–projective.*

The statement follows from the explicit formulas for tensor operators L_k and J_k.

Remark 6. For $h=0$ the $\mathcal{K}$–projective representation of the Witt algebra in the Verma module over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ is $\mathcal{FR}$–projective.

2.4. $\mathcal{B}$–projective representations of the Witt algebra. Note that one of disadvantages of many $\mathfrak{A}$–projective representations, in particular, if $\mathfrak{A}$ is a certain operator class $\mathcal{K}$, $\mathcal{HS}$, $\mathcal{TC}$ or $\mathcal{FR}$, is that their set is not closed under tensor products, in general. For the class $\mathcal{B}$ of bounded operators the tensor products of $\mathcal{B}$–projective representations are defined. However, any $\mathfrak{S}$–projective representation, where $\mathfrak{S}$ is one of classes above, is $\mathcal{B}$–projective and as such is sometimes nontrivial (when the initial representation is realized by unbounded operators), that allows to construct their tensor products, which will be nontrivial $\mathcal{B}$–projective representations in this case.

Let us call a $\mathfrak{S}$–projective representation of the Lie algebra $\mathfrak{g}$ ($\mathfrak{S}$ is a certain operator class) $\mathfrak{S}$*–irreducible* iff the representation operators can not be simultaneously transform by an addition of elements of $\mathfrak{S}$ to the form, in which they have a common proper invariant subspace.

Theorem 2. *Let T_h be the $\mathcal{B}$–projective representation of the Witt algebra $\mathfrak{w}^{\mathbb{C}}$ by the q_R–conformal symmetries in the Verma module V_h over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ then the $\mathcal{B}$–projective representations $S^n(T_h)$ (here S^n denotes the functor of the n-th symmetric degree, cf.[26]) are $\mathcal{B}$–irreducible.*

The statement of the theorem follows from the explicit formulas for generators of q_R–conformal symmetries.

Remark 7. Decompositions of the tensor products $T_{h_1}\otimes\ldots\otimes T_{h_n}$ of $\mathcal{B}$–projective representations T_{h_i} on $\mathcal{B}$–irreducible components are not known in general case.

§3. Asymptotic representations of the Witt algebra and the Virasoro algebra in Verma modules over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$

3.1. The asymptotics of q_R–conformal symmetries in Verma modules V_h over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ at $q_R\to 0$ ($h\to\frac{1}{2}$) and at $q_R\to 1$ ($h\to 1$) and the Witt algebra $\mathfrak{w}^{\mathbb{C}}$. Let $\mathfrak{A}$ be the algebra of all finite linear combinations of expressions of the form $f(qp)p^n$ ($n\in\mathbb{Z}_+$) or $q^nf(qp)$ ($n\in\mathbb{Z}_+$), where $[p,q]=1$ and f are rational functions, whose denominators have no zeroes in integer points. Let us call the algebra $\mathfrak{A}$ *the extended Weyl algebra.* The extended Weyl algebra is a topological algebra with respect to natural convergence.

Lemma. *The extended Weyl algebra $\mathfrak{A}$ admits a strict representation in the Verma module V_h ($h\notin\mathbb{Z}/2$):*

$$p\to\partial_z,\qquad q\to z.$$

Generators of q_R–conformal symmetries L_k ($k\in\mathbb{Z}$) as well as generators D, F of the Lobachevskii-Berezin algebra belong to the image of the extended Weyl algebra $\mathfrak{A}$.

Below we shall interpret the "O–large" for the asymptotics of operators in the Verma modules over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ with non-half-integer extremal weights, which belong to the extended Weyl algebra (because its representations in the Verma modules are strict one can identify the algebra itself with its image) in the sense of topology in the extended Weyl algebra.

Theorem 3A. *Generators of q_R–conformal symmetries in the Verma modules $V_{\frac{1}{2}+\hbar}$ ($0<\hbar<\frac{1}{2}$) over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ form an asymptotic "mod $O(\hbar)$" representation of the Witt algebra $\mathfrak{w}^{\mathbb{C}}$.*

Generators of q_R–conformal symmetries in the Verma modules $V_{1\pm\hbar}$ ($0<\hbar<\frac{1}{2}$) over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ form an asymptotic "mod $O(\hbar)$" representation of the Witt algebra $\mathfrak{w}^{\mathbb{C}}$, too.

The first statement of the theorem, which immediately follows from the explicit form of generators of q_R–conformal symmetries, was in fact proved in [10], where the corresponding "exponentiated" version was considered. The second is similar to the first.

Remark 8. The statement of the theorem is not true if the "O–large" is considered in sense of (weak) operator convergence in the space of operators in the Verma modules V_h over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ identified with the space $\mathbb{C}[z]$ of polynomials.

An answer on the question on a "divergence" between theorem 3A and the remark 8 is in the following: the strict representations of the extended Weyl algebra do not even exist for the limit value $\hbar = 0$, but being continuous (in a weak operator topology in the space V_h) for all values of the parameter $\hbar$ from an interval $0 < \hbar < \frac{1}{2}$, are not *uniformly* continuous on this interval. Note that the continuous representations of the extended Weyl algebra in the verma modules V_h over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ can be extended to representations by unbounded operators in the spaces V_h^{Hilb} (or V_h^{Pontr}), and, therefore, uncontinuous, that, however, is typical for the representation theory of Lie algebras.

3.2. The estimation "mod $\mathcal{HS}$" of the second term of the asymptotics of q_R–conformal symmetries in Verma modules V_h over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ at $q_R \to 0$ $(h \to \frac{1}{2})$ and at $q_R \to 1$ $(h \to 1)$ and the Virasoro algebra $\mathfrak{vir}^{\mathbb{C}}$. Under the "hybridization" of the approximations "mod $O(\hbar^n)$" and "mod $\mathfrak{S}$" an interesting phenomenon of "noncommutativity" of estimations is observed, the asymptotics of q_R–conformal symmetries supply us by an example.

Theorem 3B. *For generators of q_R–conformal symmetries in the Verma modules $V_{\frac{1}{2}+\hbar}$ $(0 < \hbar < \frac{1}{2})$ over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ the following asymptotics "mod $O(\hbar^2)$" exists:*

$$[L_i, L_j] = (i-j)L_{i+j} + \hbar\mathcal{A} + O(\hbar^2),$$

where

$$\mathcal{A} \equiv \frac{2}{3}(i^3 - i)\delta_{ij} \mod \mathcal{HS}\,.$$

So generators of q_R–conformal symmetries form an approximate representation of the Virasoro algebra $\mathfrak{vir}^{\mathbb{C}}$ for $h \to \frac{1}{2}$ in the specified sense.

For generators of q_R–conformal symmetries in the Verma modules $V_{1\pm\hbar}$ $(0<\hbar<\frac{1}{2})$ over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ the following asymptotics "mod $O(\hbar^2)$" exists, too:

$$[L_i, L_j] = (i-j)L_{i+j} + \hbar\mathcal{A} + O(\hbar^2),$$

where

$$\mathcal{A} \equiv \frac{2}{3}(i^3 - i)\delta_{ij} \mod \mathcal{HS}\,.$$

So generators of q_R–conformal symmetries form an approximate representation of the Virasoro algebra $\mathfrak{vir}^{\mathbb{C}}$ for $h \to 1$ in the specified sense.

In the statement of the theorem the estimations "mod $O(\hbar^2)$" are interpreted in the sense of a convergence in the extended Weyl algebra $\mathfrak{A}$, and $\mathcal{HS}$ denotes the set of all elements of this algebra, which are realized by Hilbert-Schmidt operators under the strict representation in the Verma modules over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ for all sufficiently small values of $\hbar$ (otherwords, if π_h is the representation of $\mathfrak{A}$ in V_h and $\mathcal{HS}_h$ is the space of Hilbert-Schmidt operators in V_h then in the theorem $\mathcal{HS}$ is interpreted as

$$\lim_{\hbar\to 0} \bigcap_{h_0<h<h_0+\hbar} \pi_h^{-1}(\mathcal{HS}_h \cap \pi_h(\mathfrak{A})),$$

where $h_0 = \frac{1}{2}$ or $h_0 = 1$).

The result of Theorem 3B follows from explicit computations. Let us compute the commutator $[L_2, L_{-2}]$ and estimate it that is enough for the determination of the central charge of the Virasoro algebra. The Verma module V_h is realized in the space $\mathbb{C}[z]$ of polynomials, $\xi = z\partial_z$, and generators of q_R–conformal symmetries are defined by expressions written above. Then for $h \to \frac{1}{2}$

$$\begin{aligned}[L_2, L_{-2}] =& \tfrac{(\xi+3h)^2(\xi+1)(\xi+2)}{(\xi+2h)(\xi+2h+1)} - \tfrac{(\xi+3h-2)^2\xi(\xi-1)}{(\xi+2h-1)(\xi+2h-2)} = \\ & \tfrac{(\xi+\frac{3}{2}+3\hbar)^2(\xi+1)(\xi+2)}{(\xi+1+2\hbar)(\xi+2+2\hbar)} - \tfrac{(\xi-\frac{1}{2}+3\hbar)^2\xi(\xi-1)}{(\xi-1+2\hbar)(\xi+2\hbar)}.\end{aligned}$$

The following estimation "mod $O(\hbar^2)$" holds ($\hbar$ in denominators is not excluded because for the zero value of $\hbar$ a representation of the extended Weyl algebra in the corresponding Verma module is not defined):

$$\begin{aligned}[L_2, L_{-2}] \equiv& (\xi + \tfrac{3}{2} + 3\hbar)^2(1 - \tfrac{2\hbar}{\xi+1+\hbar})(1 - \tfrac{2\hbar}{\xi+2+\hbar}) - \\ & (\xi - \tfrac{1}{2} + 3\hbar)^2(1 - \tfrac{2\hbar}{\xi-1+\hbar})(1 - \tfrac{2\hbar}{\xi+\hbar}) \mod O(\hbar^2).\end{aligned}$$

Extracting the terms of order $\hbar$ (because the first term of the order 1 is known) one receives:

$$2\hbar(\xi - \tfrac{1}{2})^2(\tfrac{1}{\xi-1+\hbar} + \tfrac{1}{\xi+\hbar}) - 2\hbar(\xi + \tfrac{3}{2})^2(\tfrac{1}{\xi+1+\hbar} + \tfrac{1}{\xi+2+\hbar}) + 12\hbar.$$

Perform the estimation "mod $\mathcal{HS}$" and obtain:

$$\begin{aligned}& 2\hbar\xi^2(\tfrac{1}{\xi-1+\hbar} + \tfrac{1}{\xi+\hbar} - \tfrac{1}{\xi+1+\hbar} - \tfrac{1}{\xi+2+\hbar}) - 12\hbar - \\ & 2\hbar\xi(\tfrac{1}{\xi-1-\hbar} + \tfrac{1}{\xi+\hbar}) - 6\hbar(\tfrac{1}{\xi+1+\hbar} + \tfrac{1}{\xi+2+\hbar}) \sim \\ & 2\hbar\xi^2(\tfrac{2}{(\xi-1+\hbar)(\xi+1+\hbar)} + \tfrac{2}{(\xi+\hbar)(\xi+2+\hbar)}) + 12\hbar - 16\hbar \sim \\ & 8\hbar + 12\hbar - 16\hbar = 4\hbar.\end{aligned}$$

So the "asymptotic" central charge is equal to $8\hbar$.

Analogously, for $h \to 1$

$$\begin{aligned}[L_2, L_{-2}] =& \tfrac{(\xi+3h)^2(\xi+1)(\xi+2)}{(\xi+2h)(\xi+2h+1)} - \tfrac{(\xi+3h-2)^2\xi(\xi-1)}{(\xi+2h-1)(\xi+2h-2)} = \\ & \tfrac{(\xi+3+3\hbar)^2(\xi+1)(\xi+2)}{(\xi+2+2\hbar)(\xi+3+2\hbar)} - \tfrac{(\xi+1+3\hbar)^2\xi(\xi-1)}{(\xi+2\hbar)(\xi+1+2\hbar)}.\end{aligned}$$

The following estimation "mod $O(\hbar^2)$" holds ($\hbar$ in denominators is not excluded because for the zero value of $\hbar$ a representation of the extended Weyl algebra in the corresponding Verma module is not defined):

$$\begin{aligned}[L_2, L_{-2}] \equiv& (\xi + 1)(\xi + 3 + 3\hbar)(1 - \tfrac{2\hbar}{\xi+2+\hbar})(1 - \tfrac{2\hbar}{\xi+3+\hbar})(1 + \tfrac{3\hbar}{\xi+3+\hbar}) - \\ & (\xi - 1)(\xi + 1 + 3\hbar)(1 - \tfrac{2\hbar}{\xi+\hbar})(1 - \tfrac{2\hbar}{\xi+1+\hbar})(1 + \tfrac{3\hbar}{\xi+1+\hbar}) \mod O(\hbar^2).\end{aligned}$$

Extracting the terms of order $\hbar$ (because the first term of the order 1 is known) one receives:

$$\hbar(\xi + 1)(\xi + 3)(\tfrac{1}{\xi+3+\hbar} - \tfrac{2}{\xi+2+\hbar}) - \hbar(\xi + 1)(\xi - 1)(\tfrac{1}{\xi+1+\hbar} - \tfrac{2}{\xi+\hbar}) + 6\hbar.$$

Perform the estimation "mod $\mathcal{HS}$" and obtain:

$$8\hbar + 2(\xi + 1)(\tfrac{\xi-1}{\xi+\hbar} - \tfrac{\xi+3}{\xi+2+\hbar}) \sim 8\hbar - 4\hbar = 4\hbar.$$

So the "asymptotic" central charge is equal to $8\hbar$.

Remark 9. The estimation "mod $O(\hbar^2)$" may be strengthened to "mod $O(\hbar^n)$" for any *finite* n (but not to "mod $O(\hbar^\infty)$" in view of the further estimation "mod $\mathcal{HS}$").

Remark 10. As follows from the results of the second paragraph the change of an order of estimations gives the zero central charge of the Virasoro algebra.

Perhaps, the noncommutativity of an order of the estimations "mod $O(\hbar^n)$" and "mod $\mathcal{HS}$" in the case of q_R–conformal symmetries is a partial case of a more general and fundamental fact of deviations between asymptotical theory of pseudodifferential operators and pseudodifferential calculus on asymptotic manifolds in sense of [27].

Note that the result of Theorem 3B contains a strange coincidence of "asymptotic" central charges of the Virasoro algebra for $h \to \frac{1}{2}$ and $h \to 1$ (namely, $c = 8\hbar$, $\hbar \to 0$). Perhaps, this coincidence may be explaned by the hypothetical suggestion that for all values of the extremal weight h the Virasoro algebra is presented on a certain more hidden and unexplicated level universally (i.e. its characteristics do not depend on h).

Conclusion

In the paper there are investigated various approximate representations of the infinite dimensional $\mathbb{Z}$–graded Lie algebras: the Witt algebra of all Laurent polynomial vector fields on a circle and its one-dimensional nontrivial central extension, the Virasoro algebra, by the infinite dimensional hidden symmetries in the Verma modules over the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$. There are considered as asymptotic representations "mod $O(\hbar^n)$" and representations up to a class $\mathfrak{S}$ of operators (compact operators, Hilbert-Schmidt operators and finite-rank operators) as cases, which combine both types of approximations (and in these cases an effect of noncommutativity of the order of their realizations is explicated, that perhaps is underlied by a more general fundamental fact of deviations between the asymptotical theory of pseudodifferential operators and the pseudodifferential calculus on the asymptotic manifolds in sense of [27]). Some applications of the discussed problems to the applied problems of information technologies (an organization of the information transmission in integrated videocognitive interactive systems for accelerated nonverbal computer and telecommunications) were investigated in the report [28].

References

[1] Taylor M., Pseudo-differential operators. B., 1974.

[2] Treves F., Introduction to pseudo-differential and Fourier integral operators. New York, 1980.

[3] Bogolyubov N.N., Mitropol'skiĭ Yu.A., Asymptotical methods in the theory of nonlinear oscillations [in Russian]. Moscow, Fizmatlit, 1958.

[4] Maslov V.P., Theory of perturbations and asymptotical methods [in Russian]. Moscow, Izd-vo MGU, 1965.

[5] Maslov V.P., Fedoryuk M.V., Quasiclassical approximation for equations of quantum mechanics [in Russian]. Moscow, Nauka, 1976.

[6] Karasev M.V., Maslov V.P., Nonlinear Poisson brackets. Geometry and quantization. AMS, Providence, 1993.

[7] Kirillov A.A., Elements of representation theory. Springer, 1976.

[8] Barut A., Raczka R., Theory of group representations and applications. PWN – Polish Sci. Publ. Warszawa, 1977.

[9] Juriev D., Topics in hidden symmetries. V. E-print: funct-an/9611003 (1996).

[10] Juriev D., On the infinite dimensional hidden symmetries. III. q_R–conformal symmetries at $q_R \to \infty$ and Berezin-Karasev-Maslov asymptotic quantization of $C^\infty(S^1)$. E-print: funct-an/9702002 (1997).

[11] Zhelobenko D.P., Representations of the reductive Lie algebras [in Russian]. Moscow, Nauka, 1993.

[12] Dixmier J., Algèbres enveloppantes. Gauthier-Villars, Paris, 1974.

[13] Juriev D.V., Complex projective geometry and quantum projective field theory. Theor. Math.Phys. 101(3) (1994) 1387-1403.

[14] Berezin F.A., Quantization in complex symmetric spaces [in Russian]. Izvestiya AN SSSR. Ser.matem. 39(2) (1975) 363-402.

[15] Klimek S., Lesniewski A., Quantum Riemann surfaces. I. Commun.Math.Phys. 146 (1992) 103-122.

[16] Biedenharn L., Louck J., Angular momentum in quantum mechanics. Theory and applications. Encycl.Math.Appl. V.8. Addison Wesley Publ. Comp. 1981.

[17] Fuchs D.B., Cohomology of infinite-dimensional Lie algebras [in Russian]. Moscow, Nauka, 1984.

[18] Juriev D., Noncommutative geometry, chiral anomaly in the quantum projective ($\mathfrak{sl}(2,\mathbb{C})$–invariant) field theory and $\mathfrak{jl}(2,\mathbb{C})$–invariance. J.Math.Phys. 33 (1992) 2819-2822, (E): 34 (1993) 1615.

[19] Juriev D., Remarks on nonassociative structures in quantum projective field theory: the central extension $\mathfrak{jl}(2,\mathbb{C})$ of the double $\mathfrak{sl}(2,\mathbb{C})+\mathfrak{sl}(2,\mathbb{C})$ of the simple Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ and related topics. Acta Appl.Math. 50 (1998) 191-196.

[20] Juriev D.V., Quantum projective field theory: quantum-field analogs of the Euler-Arnol'd equations in projective G-hypermultiplets. Theor.Math.Phys. 98(2) (1994) 147-161.

[21] Mal'tsev A.I., On a representation of nonassociative rings [in Russian]. Uspekhi Matem. Nauk 7(1) (1952) 181-185 [reprinted in "Selected papers. 1. Classical algebra", Moscow, Nauka, 1976, pp.328-331].

[22] Kobayashi Sh, Nomizu K., Foundations of differential geometry. Interscience Publ., New York, 1963/69.

[23] Juriev D., Infinite dimensional geometry and quantum field theory of strings. II. Infinite-dimensional noncommutative geometry of a self-interacting string field. Russian J.Math. Phys. 4 (1996) 287-314.

[24] Juriev D., On the infinite dimensional hidden symmetries. I. Infinite dimensional geometry of q_R–conformal symmetries. E-print: funct-an/9612004 (1996).

[25] Juriev D., On the infinite dimensional hidden symmetries. II. q_R–conformal modular functor. E-print: funct-an/9701009 (1997).

[26] Kirillov A.A., Introduction into the representation theory and noncommutative harmonic analysis [in Russian]. Current Math. Problems. Basic Directions. V.22, Moscow, VINITI, 1987, pp.5-162.

[27] Dubnov V.L., Maslov V.P., Nazaĭkinskiĭ V.E., The complex Lagrangian germ and the canonical operator. Russian J.Math.Phys. 3 (1995) 141-180.

[28] Juriev D.V., Droems: experimental mathematics, informatics and infinite-dimensional geometry [in Russian]. Report RCMPI-96/05$^+$ (1996).

APPENDIX: THE ORIGINAL RUSSIAN VERSION OF ARTICLE

УДК: 517.986.5+512.554.32+512.813.4

ПРИБЛИЖЕННЫЕ ПРЕДСТАВЛЕНИЯ И АЛГЕБРА ВИРАСОРО

Д.В.Юрьев

Центр математическоё физики и информатики “Таласса Этерия”,
ул.Миклухо-Маклая 20-180, Москва 117437 Россия.
E-mail: denis@juriev.msk.ru

Февраль, 1997 math.RT/9805001

Данная работа, адресованная, в основном, специалистам в теории представлениё конечномерных редуктивных и бесконечномерных $\mathbb{Z}$–градуированных алгебр Ли, теории приближениё в функциональном анализе и асимптотическим методам в исчислении операторов, а также, отчасти, в математическоё физике (алгебраическоё квантовоё теории поля), посвящена изучению алгебраическоё и аналитическоё структуры бесконечномерных скрытых симметриё в теории представлениё конечномерных редуктивных алгебр Ли. Более точно, предметом работы является реализация бесконечномерноё $\mathbb{Z}$–градуированноё алгебры Витта лорановских полиномиальных векторных полеё на окружности и ее одномерного нетривиального центрального расширения (алгебры Вирасоро) скрытыми симметриями в модулях Верма над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ (т.н. q_R–конформными симметриями). При этом, указанные бесконечномерные алгебры Ли реализуются q_R–конформными симметриями не точно, а приближенно. В статье рассматриваются два типа приближенных представлениё: представления по модулю некоторого класса $\mathfrak{S}$ операторов (операторов Гильберта-Шмидта, компактных операторов или операторов конечного ранга) и асимптотические представления “mod $O(\hbar)$”, где $\hbar$ – некоторыё параметр (в данном случае возможны различные определения операторного “O–большого” по параметру $\hbar$). Приближенные представления первого типа естественно трактовать в контексте псевдодифференциального исчисления [1,2], в то время как вторые – асимптотических методов [3-5]. Отметим, что асимптотические представления “mod $O(\hbar)$” исследовались в рамках формализма асимптотического квантования Карасева-Маслова [6], а представления по модулю $\mathfrak{S}$ являются в определенном смысле обобщениями проективных представлениё [7,8].

Результаты работы, некоторые из которых были анонсированы в электронных препринтах электронного архива по функциональному анализу Национальных лабораториё США в Лос Аламосе [9,10], носят скорее характер наблюдениё, нежели теорем, требующих трудоемких и не ясных из их формулировок доказательств. Доказательства представленных в статье фактов по сути дела одноходовы и представляют собоё громоздкие вычислительные упражнения (где это возможно, опущенные), опирающиеся на совокупность полученных автором ранее явных формул; таким образом, смысловая нагрузка приходится, в основном, на сами формулировки результатов.

§1. Предварительные определения

1.1. Алгебра Ли $\mathfrak{sl}(2,\mathbb{C})$ и модули Верма над неё. Алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ называется трехмерное пространство комплексных матриц 2×2 со следом нуль, снабженное стандартным коммутатором $[X,Y]=XY-YX$, где умножение в правоё части – стандартное умножение матриц. В базисе

$$l_{-1}=\begin{pmatrix}0&1\\0&0\end{pmatrix},\quad l_0=\begin{pmatrix}\frac{1}{2}&0\\0&-\frac{1}{2}\end{pmatrix},\quad l_1=\begin{pmatrix}0&0\\-1&0\end{pmatrix}$$

коммутационные соотношения имеют вид: $[l_i,l_j]=(i-j)l_{i+j}$ $(i,j=-1,0,1)$. Алгебра Ли $\mathfrak{sl}(2,\mathbb{C})$ $\mathbb{Z}$–градуирована: $\deg(l_i)=-\operatorname{ad}(l_0)l_i=i$, где $\operatorname{ad}(X)$ – оператор присоединенного деёствия в алгебре Ли: $\operatorname{ad}(X)Y=[X,Y]$. Как следствие, $\mathbb{Z}$–градуированные модули над $\mathfrak{sl}(2,\mathbb{C})$ являются l_0–диагональными. Вектор v в $\mathbb{Z}$–градуированном модуле над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ называется экстремальным, если $l_1v=0$ и линеёная оболочка векторов $l_{-1}^n v$ $(n\in\mathbb{Z}_+)$ совпадает с самим модулем (т.е. v – циклическиё вектор). $\mathbb{Z}$–градуированныё модуль в котором существует экстремальныё вектор (в этом случае он определен с точностью до умножения на константу) называется экстремальным [11]. Экстремальныё модуль называется модулем Верма [12], если деёствие l_{-1} в нем свободно, т.е. вектора $l_{-1}^n v$ линеёно независимы. В случае алгебры Ли $\mathfrak{sl}(2,\mathbb{C})$ модули Верма суть в точности бесконечномерные экстремальные модули. Экстремальным весом модуля Верма называется число h, определяемое из равенства $l_0v=hv$, где v – экстремальныё вектор. Модули Верма определены для всех комплексных чисел h и попарно неизоморфны. В дальнеёшем мы будем рассматривать модули Верма только с вещественными экстремальными весами.

Модуль Верма V_h над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ с экстремальным весом h может быть реализован в пространстве $\mathbb{C}[z]$ многочленов одного комплексного переменного z. Формулы для генераторов алгебры Ли $\mathfrak{sl}(2,\mathbb{C})$ имеют вид:

$$l_{-1}=z,\quad l_0=z\partial_z+h,\quad l_1=z\partial_z^2+2h\partial_z,$$

здесь $\partial_z=\frac{d}{dz}$.

Модуль Верма невырожден (т.е. не содержит собственного подмодуля) при $h\neq-\frac{n}{2}$ $(n\in\mathbb{Z}_+)$. Модуль Верма V_h называется унитаризуемым (или эрмитовым), если в нем задана структура предгильбертова пространства такая, что $l_i^*=l_{-i}$. Пополнение унитаризуемого модуля Верма будет обозначаться V_h^{Hilb}. Алгебра Ли $\mathfrak{sl}(2,\mathbb{C})$ деёствует в V_h^{Hilb} неограниченными операторами. Полезно также рассматривать формальные модули Верма V_h^{form}, которые реализуются в пространстве $\mathbb{C}[[z]]$ формальных степенных рядов одного комплексного переменного z, а формулы для генераторов алгебры Ли $\mathfrak{sl}(2,\mathbb{C})$ совпадают с приведенными выше. При этом, $V_h\subseteq V_h^{\mathrm{Hilb}}\subseteq V_h^{\mathrm{form}}$, и модули V_h, V_h^{Hilb}, V_h^{form} образуют троёку Гельфанда или оснащение гильбертова пространства V_h^{Hilb} по Дираку. Деёствие вещественноё формы алгебры Ли $\mathfrak{sl}(2,\mathbb{C})$, порожденноё антиэрмитовыми операторами il_0, $l_1 - l_{-1}$ и $i(l_1 + l_{-1})$ в гильбертовом пространстве V_h^{Hilb} неограниченными операторами, экспоненцируется до унитарного представления соответствующеё односвязноё группы Ли.

В неунитаризуемом модуле Верма над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ имеется единственная (с точностью до множителя) индефинитная полуторалинеёная форма $(\cdot,\cdot)$ такая, что $(l_i v_1, v_2) = (v_1, l_{-i} v_2)$ для любых двух векторов v_1 и v_2 из модуля Верма. Если эта полуторалинеёная форма невырождена (а в этом случае и сам модуль невырожден), то она имеет сигнатуру вида (n,∞), где n – конечно, а следовательно, определено пополнение такого модуля Верма по Понтрягину [10]. Соответствующиё модуль, в котором алгебра Ли $\mathfrak{sl}(2,\mathbb{C})$ деёствует неограниченными операторами, будет обозначаться V_h^{Pontr}. Имеет место цепочка включениё: $V_h \subseteq V_h^{\mathrm{Pontr}} \subseteq V_h^{\mathrm{form}}$. Деёствиё вещественноё формы алгебры Ли $\mathfrak{sl}(2,\mathbb{C})$, порожденноё антиэрмитовыми (относительно невырожденноё индефинитноё полуторалинеёноё формы $(\cdot,\cdot)$) операторами il_0, $l_1 - l_{-1}$ и $i(l_1 + l_{-1})$ в пространстве Понтрягина V_h^{Pontr} неограниченными операторами, экспоненцируется до псевдоунитарного представления соответствующеё односвязноё группы Ли.

1.2. Скрытые симметрии в модулях Верма над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$: C^*–алгебра Лобачевского-Березина и q_R–конформные симметрии.

Предложение 1 [13]. *В невырожденном модуле Верма V_h над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ однозначно определены операторы D и F такие, что*

$$[D, l_{-1}] = 1, \quad [D, l_0] = D, \quad [D, l_1] = D^2,$$
$$[l_{-1}, F] = 1, \quad [l_0, F] = F, \quad [l_1, F] = F^2.$$

Если модули Верма реализованы в пространстве $\mathbb{C}[z]$ многочленов одноё комплексноё переменноё z, то

$$D = \partial_z, \qquad F = z\tfrac{1}{\xi + 2h},$$

где $\xi = z\partial_z$. Операторы F и D удовлетворяют соотношениям:

$$[FD, DF] = 0, \qquad [D, F] = q_R(1 - DF)(1 - FD),$$

где $q_R = \frac{1}{2h-1}$. В унитаризуемом модуле Верма (при $q_R \neq 0$) операторы F и D ограничены и $F^* = D$, $D^* = F$.

Алгебра, порожденная переменными t и t^*, с соотношениями $[tt^*, t^*t] = 0$ и $[t, t^*] = q_R(1 - tt^*)(1 - t^*t)$, будучи квантованием по Березину плоскости Лобачевского, реализованноё в единичном комплексном диске (реализация Пуанкаре) [14], называется алгеброё Лобачевского-Березина. Предложение 1 позволяет рассматривать алгебру Лобачевского-Березина как C^*–алгебру. C^*–алгебра Лобачевского-Березина была недавно переоткрыта С.Климеком и А.Лесниевским [15].

Предложение 2 [13]. *В невырожденном модуле Верма V_h над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ однозначно определены операторы L_n ($n \in \mathbb{Z}$) такие, что*

$$[l_i, L_n] = (i-n)L_{i+n} \qquad (i = 1, 2, 1;\ n \in \mathbb{Z}),$$

при этом, $L_i = l_i$ ($i = -1, 0, 1$). Если модули Верма реализованы в пространстве $\mathbb{C}[z]$ многочленов одноё комплексноё переменноё z, то

$$L_k = (xi + (k+1)h)\partial_z^k\ (k \geq 0), \quad L_{-k} = z^k \tfrac{\xi + (k+1)h}{(\xi+2h)\ldots(\xi+2h+k-1)}\ (k \geq 1),$$

где $\xi = z\partial_z$. Операторы L_n удовлетворяют соотношениям:

$$[L_n, L_m] = (n-m)L_{n+m},\ \text{если}\ n, m \geq -1\ \text{или}\ n, m \leq 1.$$

В унитаризуемом модуле Верма операторы L_n неограничены и $L_i^* = L_{-1}$.

Операторы L_n называются q_R–конформными симметриями. Они *символически* могут быть представлены в виде:

$$L_n = D^{nh} L_0 D^{n(1-h)}, \qquad L_{-n} = F^{n(1-h)} L_0 F^{nh}.$$

Для придания смысла символическоё записи следует воспользоваться общими перестановочными соотношениями

$$[L_n, f(D)] = (-D)^{n+1} f'(D)\ (n \geq -1), \qquad [L_{-n}, f(F)] = F^{n+1} f'(D)\ (n \geq -1)$$

при $n = 0$.

Соотношения коммутации операторов D, F и генераторов q_R–конформных симметриё с генераторами алгебры Ли $\mathfrak{sl}(2,\mathbb{C})$ означают, что семеёства J_k и L_k ($k \in \mathbb{Z}$), где $J_i = D^i$, $J_{-i} = F^i$ ($i \in \mathbb{Z}_+$), являются семеёствами тензорных операторов [8,16] для алгебры Ли $\mathfrak{sl}(2,\mathbb{C})$.

С семеёствами тензорных операторов J_k и L_k можно ассоциировать их производящие функции – операторные поля, т.е. формальные ряды Лорана от одноё комплексноё переменноё u с операторными коэффициентами:

$$J(u) = \sum_{i \in \mathbb{Z}} J_i (-u)^{-1-i}, \quad T(u) = \sum_{i \in \mathbb{Z}} L_i (-u)^{-2-i}.$$

При каждом значении u операторное поле задает отображение V_h в V_h^{form}. Тот факт, что J_k и T_k ($k \in \mathbb{Z}$) образуют семеёства тензорных операторов, на языке операторных полеё означает, что

$$[l_i, J(u)] = (-u)^i J(u) - (-u)^{i+1} J'(u), \quad [l_i, T(u)] = 2(-u)^i T(u) - (-u)^{i+1} T'(u).$$

Операторные поля $V(u)$, преобразующиеся как s–дифференциалы под деёствием алгебры Ли $\mathfrak{sl}(2,\mathbb{C})$ (т.е. удовлетворяющие тождеству $[l_i, V(u)] = s(-u)^i V(u) - (-u)^{i+1} V'(u)$), называются $\mathfrak{sl}(2,\mathbb{C})$–первичными спина s; таким образом, операторные поля $J(u)$ и $T(u)$ являются $\mathfrak{sl}(2,\mathbb{C})$–первичными спина 1 и 2, соответственно. Операторные поля $J(u)$ и $T(u)$, а также их своёства подробно изучались в [13].

1.3. Бесконечномерные $\mathbb{Z}$–градуированные алгебры Ли: алгебра Витта $\mathfrak{w}^{\mathbb{C}}$ лорановских полиномиальных векторных полеё на окружности и алгебра Вирасоро $\mathfrak{vir}^{\mathbb{C}}$, ее одномерное нетривиальное центральное расширение. Алгебра Ли $\mathrm{Vect}(S^1)$ реализуется в пространстве C^∞–гладких векторных полеё $v(t)\partial_t$ на окружности $S^1 \simeq \mathbb{R}/2\pi\mathbb{Z}$ с коммутатором

$$[v_1(t)\partial_t, v_2(t)\partial_t] = (v_1(t)v_2'(t) - v_1'(t)v_2(t))\partial_t.$$

В базисе $s_n = \sin(nt)\partial_t$, $c_n = \cos(nt)\partial_t$, $h = \partial_t$ коммутационные соотношения имеют вид:

$$\begin{aligned}
[s_n, s_m] &= \tfrac{1}{2}((m-n)s_{n+m} + \mathrm{sgn}(n-m)(n+m)s_{|n-m|}),\\
[c_n, c_m] &= \tfrac{1}{2}((n-m)s_{n+m} + \mathrm{sgn}(n-m)(n+m)s_{|n-m|}),\\
[s_n, c_m] &= \tfrac{1}{2}((m-n)c_{n+m} - (m+n)c_{|n-m|}) - n\delta_{nm}h,\\
[h, s_n] &= nc_n, \quad [h, c_n] = -ns_n.
\end{aligned}$$

Обозначим $\mathrm{Vect}^{\mathbb{C}}(S^1)$ комплексификацию алгебры Ли $\mathrm{Vect}(S^1)$. В базисе $e_n = ie^{ikt}\partial_t$ коммутационные соотношения в алгебре Ли $\mathrm{Vect}^{\mathbb{C}}(S^1)$ имеют вид:

$$[e_j, e_k] = (j-k)e_{j+k}.$$

Удобно также рассматривать вложение окружности S^1 в комплексную плоскость $\mathbb{C}$ с координатоё z, при этом на окружности $z = e^{it}$, а элементы базиса e_k $(k \in \mathbb{Z})$ представляются лорановскими полиномиальными векторными полями: $e_k = z^{k+1}\partial_z$. $\mathbb{Z}$–градуированная алгебра Ли, порожденная лорановскими полиномиальными векторными полями (т.е. конечными линеёными комбинациями элементов базиса e_k), называется алгеброё Витта и обозначается $\mathfrak{w}^{\mathbb{C}}$. Алгебра Витта $\mathfrak{w}^{\mathbb{C}}$ является комплексификациеё подалгебры $\mathfrak{w}$ алгебры $\mathrm{Vect}(S^1)$, порожденноё тригонометрическими полиномиальными векторными полями на окружности S^1, т.е. конечными линеёными комбинациями элементов базиса s_n, c_n и h.

Алгебра Ли $\mathrm{Vect}(S^1)$ допускает нетривиальное одномерное центральное расширение, задаваемое 2-коциклом Гельфанда-Фукса [17]:

$$c(v_1(t)\partial_t, v_2(t)\partial_t) = \int_0^{2\pi} (v_1'(t)v_2''(t) - v_2'(t)v_1'(t))\,dt.$$

Данное расширение, будучи продолженным на комплексификацию $\mathrm{Vect}^{\mathbb{C}}(S^1)$ алгебры Ли $\mathrm{Vect}(S^1)$ и ограниченным на подалгебру $\mathfrak{w}^{\mathbb{C}}$, задает одномерное центральное расширение алгебры Витта, называемое алгеброё Вирасоро и обозначаемое $\mathfrak{vir}^{\mathbb{C}}$. Алгебра Вирасоро порождается генераторами e_k $(k \in \mathbb{Z})$ и центральным элементом c с коммутационными соотношениями:

$$[e_j, e_k] = (j-k)e_{j+k} + \frac{j^3 - j}{12}c$$

и является комплексификациеё центрального расширения $\mathfrak{vir}$ алгебры Ли $\mathfrak{w}$. В неприводимом представлении центральныё элемент c алгебры Вирасоро переходит в скалярныё оператор, коэффициент пропорциональности которого единичному оператору называется *центральным зарядом.*

§2. $\mathcal{HS}$–проективные представления алгебры Витта в модулях Верма над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$

2.1. $\mathfrak{A}$–проективные представления [9].

Определение 1А. Пусть $\mathfrak{A}$ – произвольная ассоциативная алгебра, представленная линеёными операторами в линеёном пространстве H и $\mathfrak{g}$ – алгебра Ли. Линеёное отображение $T:\mathfrak{g}\mapsto \operatorname{End}(H)$ называется $\mathfrak{A}$*–проективным представлением*, если для произвольных X и Y из $\mathfrak{g}$ существует элемент алгебры $\mathfrak{A}$, представленныё оператором A_{XY}, такоё, что

$$[T(X),T(Y)]-T([X,Y])=A_{XY}.$$

Если H бесконечномерно, то представление может осуществляться неограниченными операторами с подходящими областями определения.

Замечание 1. Определение 1А может быть обобщено на произвольные антикоммутативные алгебры. В этоё ситуации оно тесно связано с конструкциями представлениё антикоммутативных алгебр $\mathfrak{jl}(2,\mathbb{C})$ и $\mathfrak{sl}^*(2,\mathbb{C})$ в [18,19;20:§2]. В целом, его следует рассматривать в контексте старых идеё А.И.Мальцева о представлениях произвольных неассоциативных алгебр [21]. Общие антикоммутативные алгебры и конструкции их $\mathfrak{A}$–проективных представлениё интересны с точки зрения квазихопфовых алгебр, неассоциативных как коалгебры, якобиевых и коякобиевых квазибиалгебр и связанных с ними структур (см.ссылки в [9]).

Пример. Если $(\mathfrak{g},\mathfrak{h})$ – редуктивная пара, то любое представление алгебры Ли $\mathfrak{g}$ является $\mathcal{U}(\mathfrak{h})$–проективным представлением бинарноё антикоммутативноё алгебры $\mathfrak{p}$ ($\mathfrak{g}=\mathfrak{h}\oplus\mathfrak{p}$, бинарная операция в $\mathfrak{p}$ имеет стандартныё вид: $[X,Y]_{\mathfrak{p}}=\pi([X,Y])$, где $[\cdot,\cdot]$ – коммутатор в $\mathfrak{g}$, а π – проектор $\mathfrak{g}$ на $\mathfrak{p}$ вдоль $\mathfrak{h}$ [22]).

Замечание 2. Стандартное проективное представление является частным случаем определения, если алгебра $\mathfrak{A}$ одномерна и деёствует в H скалярными матрицами.

Замечание 3. Если H – гильбертово (или предгильбертово) пространство, то в качестве $\mathfrak{A}$ можно рассматривать алгебру $\mathcal{HS}$ всех операторов Гильберта-Шмидта. Можно также рассматривать алгебры $\mathcal{B}$, $\mathcal{K}$, $\mathcal{TC}$ и $\mathcal{FR}$ ограниченных, компактных, ядерных операторов и операторов конечного ранга.

Определение 1Б. Пусть $\mathfrak{A}$ – ассоциативная алгебра с инволюциеё $*$, симметрично представленная в гильбертовом пространстве H. Если $\mathfrak{g}$ – алгебра Ли с инволюциеё $*$, то ее $\mathfrak{A}$–проективное представление T в пространстве H называется *симметричным*, если для всех элементов a из $\mathfrak{g}$ $T(a^*)=T^*(a)$. Пусть $\mathfrak{g}$ – $\mathbb{Z}$–градуированная алгебра Ли ($\mathfrak{g}=\oplus_{n\in\mathbb{Z}}\mathfrak{g}_n$) с инволюциеё $*$ такая, что $\mathfrak{g}_n^*=\mathfrak{g}_{-n}$ и инволюция тождественна на подалгебре $\mathfrak{g}_0$. Продолжим $\mathbb{Z}$–градуировку и инволюцию $*$ с алгебры Ли $\mathfrak{g}$ на тензорную алгебру $\mathbf{T}^{\cdot}(\mathfrak{g})$. Симметричное $\mathfrak{A}$–проективное представление $\mathfrak{g}$ называется *абсолютно симметричным*, если для любого элемента a алгебры $\mathbf{T}^{\cdot}(\mathfrak{g})$ такого, что $\deg(a)=0$, выполнено равенство $T(a)=T^*(a)$ (здесь представление T алгебры Ли $\mathfrak{g}$ в H продолжено до отображения $\mathbf{T}^{\cdot}(\mathfrak{g})$ в $\operatorname{End}(H)$.

Определение 1B. $\mathfrak{A}$–проективное представление T алгебры Ли $\mathfrak{g}$ в линеёном пространстве H называется *почти абсолютно замкнутым*, если для любого натурального n и произвольных элементов $X_0, X_1, X_2, \dots X_{n+1}$ алгебры Ли $\mathfrak{g}$ существует элемент $\varphi(X_0, X_1, X_2, \dots X_{n+1})$ алгебры $\mathfrak{g}$ такоё, что

$$[\dots[[T(X_0), T(X_1)], T(X_2)], \dots, T(X_{n+1})] \equiv T(\varphi(X_0, X_1, X_2, \dots X_{n+1})) \pmod{\mathfrak{A}},$$

здесь $\mathfrak{A}$ рассматривается представленноё в $\mathrm{End}(H)$. Почти абсолютно замкнутое $\mathfrak{A}$–проективное представление T алгебры Ли $\mathfrak{g}$ в линеёном пространстве H называется *абсолютно замкнутым*, если $\varphi(\cdot, \dots, \cdot) \equiv 0$.

Отображения $(X_0, X_1, X_2, \dots X_{n+1}) \mapsto \varphi(X_0, X_1, X_2, \dots X_{n+1})$, ассоциированные с произвольным почти абсолютно замкнутым $\mathfrak{A}$–проективным представлением алгебры Ли $\mathfrak{g}$, определяют высшие скобки в алгебре Ли $\mathfrak{g}$. Объекты с высшими скобками систематически появляются во многих областях математики и математическоё физики (см.напр.книгу [6] или статью [23] среди прочих и многочисленные ссылки в них).

2.2. $\mathcal{HS}$–проективные представления алгебры Витта q_R–конформными симметриями в унитаризуемых модулях Верма V_h над алгеброё Ли $\mathfrak{sl}(2, \mathbb{C})$ [9]. Отметим, что алгебра Витта $\mathfrak{w}^{\mathbb{C}}$ допускает естественную инволюцию $*$.

Теорема 1A. *Генераторы L_k $(k \in \mathbb{Z})$ q_R–конформных симметриё в унитаризуемом модуле Верма V_h над алгеброё Ли $\mathfrak{sl}(2, \mathbb{C})$ осуществляют абсолютно симметричное $\mathcal{HS}$–проективное представление алгебры Витта $\mathfrak{w}^{\mathbb{C}}$. Добавление к тензорным операторам L_k тензорных операторов J_k $(k \in \mathbb{Z})$ приводит к абсолютно симметричному $\mathcal{HS}$–проективному представлению полупрямоё суммы алгебры Витта и бесконечномерноё $\mathbb{Z}$–градуированноё алгебры Геёзенберга (одномерного центрального расширения бесконечномерноё $\mathbb{Z}$–градуированноё абелевоё алгебры Ли $\mathbb{C}[z, z^{-1}]$ многочленов Лорана).*

Утверждение теоремы следует из явных формул для генераторов q_R–конформных симметриё и тензорных операторов J_k. Проверка на принадлежность классу $\mathcal{HS}$ операторов Гильберта-Шмидта для операторов фиксированноё степени относительно градуировки в $\mathbb{Z}$–градуированном пространстве многочленов $\mathbb{C}[z]$, снабженном некоторым скалярным произведением, относительно которого одномерные весовые пространства ортогональны, каковым и является унитаризуемыё модуль Верма над алгеброё Ли $\mathfrak{sl}(2, \mathbb{C})$, не представляет никаких проблем.

Замечание 4. $\mathcal{HS}$–проективные представления алгебры Витта в унитаризуемых модулях Верма над $\mathfrak{sl}(2, \mathbb{C})$ абсолютно замкнуты.

Замечание 5. Теорема 1A переносится на псевдоунитарныё случаё с заменоё класса $\mathcal{HS}$ операторов Гильберта-Шмидта на класс $\mathcal{K}$ компактных операторов.

Результаты, касающиеся “экспоненцированноё” версии теоремы 1A, анонсированы в электронных препринтах [24,25].

2.3. $\mathcal{FR}$–проективные представления алгебры Витта в модулях Верма над алгеброё Ли $\mathfrak{sl}(2, \mathbb{C})$ с экстремальным весом 1 и $\frac{1}{2}$. В некоторых частных случаях утверждение теоремы 1A может быть усилено.

Теорема 1Б. *При* $h=1$ *или* $h=\frac{1}{2}$ $\mathcal{HS}$*–проективное представление алгебры Витта* q_R*–конформными симметриями в унитаризуемых модулях Верма над алгеброӗ Ли* $\mathfrak{sl}(2,\mathbb{C})$ *является* $\mathcal{FR}$*–проективным. При* $h=\frac{1}{2}$ $\mathcal{FR}$*–проективным является и* $\mathcal{HS}$*–проективное представление полупрямоӗ суммы алгебры Витта и бесконечномерноӗ алгебры Геӗзенберга.*

Утверждение следует из явных формул для тензорных операторов L_k и J_k.

Замечание 6. При $h=0$ $\mathcal{K}$–проективное представление алгебры Витта в модуле Верма над алгеброӗ Ли $\mathfrak{sl}(2,\mathbb{C})$ является $\mathcal{FR}$–проективным.

2.4. $\mathcal{B}$–проективные представления алгебры Витта. Отметим, что к числу "неудобств" многих $\mathfrak{A}$–проективных представлениӗ, в том числе если $\mathfrak{A}$ – некоторыӗ класс операторов $\mathcal{K}$, $\mathcal{HS}$, $\mathcal{TC}$ или $\mathcal{FR}$, относится, вообще говоря, незамкнутость их совокупности относительно взятия тензорных произведениӗ. Для класса же $\mathcal{B}$ ограниченных операторов тензорные произведения $\mathcal{B}$–проективных представлениӗ определены. Однако, всякое $\mathfrak{S}$–проективное представление, где $\mathfrak{S}$ – один из упомянутых выше классов, является $\mathcal{B}$–проективным и как таковое, иногда, нетривиальным (когда исходные представления осуществлялись неограниченными операторами), что позволяет конструировать их тензорные произведения, которые будут в этом случае нетривиальными $\mathcal{B}$–проективными представлениями.

Назовем $\mathfrak{S}$–проективное представление алгебры Ли $\mathfrak{g}$ ($\mathfrak{S}$ – некоторыӗ операторныӗ класс) $\mathfrak{S}$*–неприводимым*, если операторы представления одновременно не могут быть приведены путем добавления к ним элементов из $\mathfrak{S}$ к виду, в котором они все обладали бы общим для них собственным инвариантным подпространством.

Теорема 2. *Пусть* T_h *–* $\mathcal{B}$*–проективное представление алгебры Витта* $\mathfrak{w}^{\mathbb{C}}$ q_R*–конформными симметриями в модуле Верма* V_h *над алгеброӗ Ли* $\mathfrak{sl}(2,\mathbb{C})$*, тогда* $\mathcal{B}$*–проективные представления* $S^n(T_h)$ *(здесь* S^n *обозначает функтор взятия* n*–оӗ симметрической степени, ср.[26])* $\mathcal{B}$*–неприводимы.*

Утверждение теоремы следует из явных формул для генераторов q_R–конформных симметриӗ.

Замечание 7. Разложения тензорных произведениӗ $T_{h_1}\otimes\ldots\otimes T_{h_n}$ $\mathcal{B}$–проективных представлениӗ T_{h_i} на $\mathcal{B}$–неприводимые компоненты в общем случае неизвестны.

§3. Асимптотические представления алгебр Витта и Вирасоро в модулях Верма над алгеброӗ Ли $\mathfrak{sl}(2,\mathbb{C})$

3.1. Асимптотика q_R–конформных симметриӗ в модулях Верма V_h над алгеброӗ Ли $\mathfrak{sl}(2,\mathbb{C})$ при $q_R\to 0$ ($h\to\frac{1}{2}$) и $q_R\to 1$ ($h\to 1$) и алгебра Витта $\mathfrak{w}^{\mathbb{C}}$. Пусть $\mathfrak{A}$ – алгебра конечных линеӗных комбинациӗ выражениӗ вида $f(qp)p^n$ $(n\in\mathbb{Z}_+)$ или $q^nf(qp)$ $(n\in\mathbb{Z}_+)$, $[p,q]=1$, а f – рациональные функции, чьи знаменатели не имеют нулеӗ в целых точках. Будем называть $\mathfrak{A}$ *расширенноӗ алгеброӗ Веӗля.* Расширенная алгебра Веӗля является топологическоӗ алгеброӗ относительно естественноӗ сходимости.

Лемма. *Расширенная алгебра Веёля* $\mathfrak{A}$ *допускает точное представление в модуле Верма* V_h $(h \notin \mathbb{Z}/2)$*:*

$$p \to \partial_z, \qquad q \to z.$$

При этом, генераторы q_R*–конформных симметриё* L_k $(k \in \mathbb{Z})$*, равно как и генераторы* D*,* F *алгебры Лобачевского-Березина, принадлежат образу расширенноё алгебры Веёля* $\mathfrak{A}$*.*

Будем понимать в дальнеёшем "O–большое" для асимптотик операторов в модулях Верма над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ с неполуцелыми экстремальными весами, принадлежащих расширенноё алгебре Веёля (поскольку ее представления в модулях Верма точны можно отождествлять саму алгебру с ее образом), в смысле топологии в расширенноё алгебре Веёля.

Теорема 3А. *Генераторы* q_R*–конформных симметриё в модулях Верма* $V_{\frac{1}{2}+\hbar}$ $(0<\hbar<\frac{1}{2})$ *над алгеброё Ли* $\mathfrak{sl}(2,\mathbb{C})$ *образуют асимптотическое* "mod $O(\hbar)$" *представление алгебры Витта* $\mathfrak{w}^{\mathbb{C}}$*.*

Генераторы q_R*–конформных симметриё в модулях Верма* $V_{1\pm\hbar}$ $(0<\hbar<\frac{1}{2})$ *над алгеброё Ли* $\mathfrak{sl}(2,\mathbb{C})$ *также образуют асимптотическое* "mod $O(\hbar)$" *представление алгебры Витта* $\mathfrak{w}^{\mathbb{C}}$*.*

Первое утверждение теоремы, немедленно следующее из явного вида генераторов q_R–конформных симметриё, было по сути дела доказано в [10], где рассматривалась соответствующая "экспоненцированная" версия. Второе ему полностью подобно.

Замечание 8. Утверждение теоремы перестает быть верным, елси "O–большое" рассматривается в смысле (слабоё) операторноё сходимости в пространстве операторов в модулях Верма V_h над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$, отождествленных с пространством $\mathbb{C}[z]$ полиномов.

Ответ на вопрос о причинах "расхождения" между теоремоё 3А и замечанием 8 заключается в следующем: точные представления расширенноё алгебры Веёля не только не существуют при предельном значении $\hbar=0$, но будучи непрерывными (в слабоё операторноё топологии в пространстве V_h) при всех значениях параметра $\hbar$ из интервала $0<\hbar<\frac{1}{2}$, не являются *равномерно* непрерывными на этом интервале. Отметим также, что непрерывные представления расширенноё алгебры Веёля в модулях Верма V_h над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ продолжаются до представлениё неограниченными операторами в пространствах V_h^{Hilb} (или V_h^{Pontr}), и, как следствие, не непрерывных, что, однако, типично для теории представлениё алгебр Ли.

3.2. Оценка "mod $\mathcal{HS}$" второго члена асимптотики q_R–конформных симметриё в модулях Верма V_h над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ при $q_R\to 0$ $(h\to\frac{1}{2})$ и при $q_R\to 1$ $(h\to 1)$ и алгебра Вирасоро $\mathfrak{vir}^{\mathbb{C}}$. При "гибридизации" приближениё "mod $O(\hbar^n)$" и "mod $\mathfrak{S}$" наблюдается интересное явление "некоммутируемости" оценок, пример чему предоставляют асимптотики q_R–конформных симметриё.

Теорема 3Б. *Для генераторов q_R–конформных симметриё в модулях Верма $V_{\frac{1}{2}+\hbar}$ $(0<\hbar<\frac{1}{2})$ над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ имеется следующая асимптотика “mod $O(\hbar^2)$”:*

$$[L_i, L_j] = (i-j)L_{i+j} + \hbar\mathcal{A} + O(\hbar^2),$$

где

$$\mathcal{A} \equiv \frac{2}{3}(i^3 - i)\delta_{ij} \mod \mathcal{HS}\,.$$

Таким образом, генераторы q_R–конформных симметриё образуют приближенное представление алгебры Вирасоро $\mathfrak{vir}^{\mathbb{C}}$ при $h \to \frac{1}{2}$ в указанном смысле.

Для генераторов q_R–конформных симметриё в модулях Верма $V_{1\pm\hbar}$ $(0<\hbar<\frac{1}{2})$ над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ также имеется асимптотика “mod $O(\hbar^2)$”:

$$[L_i, L_j] = (i-j)L_{i+j} + \hbar\mathcal{A} + O(\hbar^2),$$

где

$$\mathcal{A} \equiv \frac{2}{3}(i^3 - i)\delta_{ij} \mod \mathcal{HS}\,.$$

Таким образом, генераторы q_R–конформных симметриё образуют приближенное представление алгебры Вирасоро $\mathfrak{vir}^{\mathbb{C}}$ при $h \to 1$ в указанном смысле.

В утверждении теоремы оценки “mod $O(\hbar^2)$” понимаются в смысле сходимости в расширенноё алгебре Веёля $\mathfrak{A}$, при этом $\mathcal{HS}$ обозначает совокупность элементов этоё алгебры, реализуемых операторами Гильберта-Шмидта при точном представлении в модулях Верма над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$ для всех достаточно малых значениё $\hbar$ (иными словами, если π_h – представление $\mathfrak{A}$ в V_h и $\mathcal{HS}_h$ – пространство операторов Гильберта-Шмидта в V_h, то $\mathcal{HS}$ понимается в теореме в смысле

$$\lim_{\hbar\to 0} \bigcap_{h_0<h<h_0+\hbar} \pi_h^{-1}(\mathcal{HS}_h \cap \pi_h(\mathfrak{A})),$$

где $h_0 = \frac{1}{2}$ или $h_0 = 1$).

Результат теоремы 3Б следует из явных вычислениё. Приведем вычисление коммутатора $[L_2, L_{-2}]$ и его оценки, которых достаточно для определения центрального заряда алгебры Вирасоро. Модуль Верма V_h реализован в пространстве многочленов $\mathbb{C}[z]$, $\xi = z\partial_z$, а генераторы q_R–конформных симметриё задаются выражениями, выписанными ранее. Тогда при $h \to \frac{1}{2}$

$$\begin{aligned}[L_2, L_{-2}] = & \tfrac{(\xi+3h)^2(\xi+1)(\xi+2)}{(\xi+2h)(\xi+2h+1)} - \tfrac{(\xi+3h-2)^2\xi(\xi-1)}{(\xi+2h-1)(\xi+2h-2)} = \\ & \tfrac{(\xi+\frac{3}{2}+3\hbar)^2(\xi+1)(\xi+2)}{(\xi+1+2\hbar)(\xi+2+2\hbar)} - \tfrac{(\xi-\frac{1}{2}+3\hbar)^2\xi(\xi-1)}{(\xi-1+2\hbar)(\xi+2\hbar)}.\end{aligned}$$

Имеыет место следующая оценка “mod $O(\hbar^2)$” ($\hbar$ в знаменателях оставлен, т.к. при нулевом $\hbar$ не определено представление расширенноё алгебры Веёля в соответствующем модуле Верма):

$$\begin{aligned}[L_2, L_{-2}] \equiv & (\xi + \tfrac{3}{2} + 3\hbar)^2(1 - \tfrac{2\hbar}{\xi+1+\hbar})(1 - \tfrac{2\hbar}{\xi+2+\hbar}) - \\ & (\xi - \tfrac{1}{2} + 3\hbar)^2(1 - \tfrac{2\hbar}{\xi-1+\hbar})(1 - \tfrac{2\hbar}{\xi+\hbar}) \mod O(\hbar^2).\end{aligned}$$

Выделяя члены порядка $\hbar$ (поскольку старшиё член порядка единицы уже известен), получим:

$$2\hbar(\xi-\tfrac{1}{2})^2(\tfrac{1}{\xi-1+\hbar}+\tfrac{1}{\xi+\hbar})-2\hbar(\xi+\tfrac{3}{2})^2(\tfrac{1}{\xi+1+\hbar}+\tfrac{1}{\xi+2+\hbar})+12\hbar.$$

Проведем теперь оценку “mod $\mathcal{HS}$” и получим:

$$\begin{aligned}&2\hbar\xi^2(\tfrac{1}{\xi-1+\hbar}+\tfrac{1}{\xi+\hbar}-\tfrac{1}{\xi+1+\hbar}-\tfrac{1}{\xi+2+\hbar})-12\hbar-\\&2\hbar\xi(\tfrac{1}{\xi-1-\hbar}+\tfrac{1}{\xi+\hbar})-6\hbar(\tfrac{1}{\xi+1+\hbar}+\tfrac{1}{\xi+2+\hbar})\sim\\&2\hbar\xi^2(\tfrac{2}{(\xi-1+\hbar)(\xi+1+\hbar)}+\tfrac{2}{(\xi+\hbar)(\xi+2+\hbar)})+12\hbar-16\hbar\sim\\&8\hbar+12\hbar-16\hbar=4\hbar.\end{aligned}$$

Таким образом, “асимптотическиё” центральныё заряд равен $8\hbar$.

Аналогично, при $h\to 1$

$$\begin{aligned}[L_2,L_{-2}]=&\tfrac{(\xi+3h)^2(\xi+1)(\xi+2)}{(\xi+2h)(\xi+2h+1)}-\tfrac{(\xi+3h-2)^2\xi(\xi-1)}{(\xi+2h-1)(\xi+2h-2)}=\\&\tfrac{(\xi+3+3\hbar)^2(\xi+1)(\xi+2)}{(\xi+2+2\hbar)(\xi+3+2\hbar)}-\tfrac{(\xi+1+3\hbar)^2\xi(\xi-1)}{(\xi+2\hbar)(\xi+1+2\hbar)}.\end{aligned}$$

Имеет место следующая оценка “mod $O(\hbar^2)$” ($\hbar$ в знаменателях оставлен, т.к. при нулевом $\hbar$ не определено представление расширенноё алгебры Веёля в соответствующем модуле Верма):

$$\begin{aligned}[L_2,L_{-2}]\equiv&(\xi+1)(\xi+3+3\hbar)(1-\tfrac{2\hbar}{\xi+2+\hbar})(1-\tfrac{2\hbar}{\xi+3+\hbar})(1+\tfrac{3\hbar}{\xi+3+\hbar})-\\&(\xi-1)(\xi+1+3\hbar)(1-\tfrac{2\hbar}{\xi+\hbar})(1-\tfrac{2\hbar}{\xi+1+\hbar})(1+\tfrac{3\hbar}{\xi+1+\hbar})\mod O(\hbar^2).\end{aligned}$$

Выделяя члены порядка $\hbar$ (поскольку старшиё член порядка единицы уже известен), получим:

$$\hbar(\xi+1)(\xi+3)(\tfrac{1}{\xi+3+\hbar}-\tfrac{2}{\xi+2+\hbar})-\hbar(\xi+1)(\xi-1)(\tfrac{1}{\xi+1+\hbar}-\tfrac{2}{\xi+\hbar})+6\hbar.$$

Проведем теперь оценку “mod $\mathcal{HS}$” и получим:

$$8\hbar+2(\xi+1)(\tfrac{\xi-1}{\xi+\hbar}-\tfrac{\xi+3}{\xi+2+\hbar})\sim 8\hbar-4\hbar=4\hbar.$$

Таким образом, “асимптотическиё” центральныё заряд опять равен $8\hbar$.

Замечание 9. Оценка “mod $O(\hbar^2)$” может быть улучшена до “mod $O(\hbar^n)$” для любого *конечного* n (но не до “mod $O(\hbar^\infty)$” из-за последующеё оценки “mod $\mathcal{HS}$”).

Замечание 10. Как следует из результатов второго параграфа перестановка порядка оценок приводит к нулевому центральному заряду для алгебры Вирасоро.

По-видимому, неперестановочность порядка оценок “mod $O(\hbar^n)$” и “mod $\mathcal{HS}$” в случае q_R–конформных симметриё является отражением более общего и фундаментального факта различиё между асимптотическоё теориeё псевдодифференциальных операторов и псевдодифференциальным исчислением на асимптотических многообразиях в смысле [27].

Отметим, что результат теоремы 3Б содержит странное совпадение “асимптотических” центральных зарядов для алгебры Вирасоро при $h\to\frac{1}{2}$ и $h\to 1$ (а именно, $c=8\hbar$, $\hbar\to 0$). Возможно, это совпадение объясняется тем, что при всех значениях экстремального веса h алгебра Вирасоро присутствует на некотором еще более скрытом и пока не выявленном уровне универсально (т.е. ее характеристики не зависят от h).

Заключение

В работе исследованы различные приближенные представления бесконечномерных $\mathbb{Z}$–градуированных алгебр Ли: алгебры Витта лорановских полиномиальных векторных полеё на окружности и ее одномерного нетривиального центрального расширения, алгебры Вирасоро, бесконечномерными скрытыми симметриями в модулях Верма над алгеброё Ли $\mathfrak{sl}(2,\mathbb{C})$. Рассмотрены как асимптотические представления “mod $O(\hbar^n)$” и представления с точностью до операторов из некоторого класса $\mathfrak{S}$ (компактных операторов, операторов Гильберта-Шмидта или операторов конечного ранга), так и случаи, совмещающие оба типа приближениё (и в данном случае выявлен эффект неперестановочности порядка их выполнения, что, по-видимому, свидетельствует о более общем и фундаментальном факте различиё между асимптотическоё теориеё псевдодифференциальных операторов и псевдодифференциальным исчислением на асимптотических многообразиях в смысле [27]). Некоторые приложения обсуждаемых вопросов к прикладным проблемам информационных технологиё (организация передачи информации в интегрированных видеокогнитивных интерактивных системах для ускоренных невербальных компьютерных и телекоммуникациё) изучались в работе [28].

Список литературы

[1] Тэёлор М., Псевдодифференциальные операторы. М., Мир, 1985.

[2] Трев Ф., Введение в псевдодифференциальные операторы и интегральные операторы Фурье. М., Мир, 1984.

[3] Боголюбов Н.Н., Митропольскиё Ю.А., Асимптотические методы в теории нелинеёных колебаниё. М., Физматлит, 1958.

[4] Маслов В.П., Теория возмущениё и асимптотические методы. М., Изд-во МГУ, 1965.

[5] Маслов В.П., Федорюк М.В., Квазиклассическое приближение для уравнениё квантовоё механики. М., Наука, 1976.

[6] Карасев М.В., Маслов В.П., Нелинеёные скобки Пуассона. Геометрия и квантование. М., Наука, 1991.

[7] Кириллов А.А., Элементы теории представлениё. М., Наука, 1978.

[8] Барут А., Рончка Р., Теория представлениё групп и ее приложения. М., Мир, 1980.

[9] Juriev D., Topics in hidden symmetries. V. E-print: funct-an/9611003 (1996).

[10] Juriev D., On the infinite dimensional hidden symmetries. III. q_R–conformal symmetries at $q_R \to \infty$ and Berezin-Karasev-Maslov asymptotic quantization of $C^\infty(S^1)$. E-print: funct-an/9702002 (1997).

[11] Желобенко Д.П., Представления редуктивных алгебр Ли. М., Наука, 1993.

[12] Диксмье Ж., Универсальные обертывающие алгебры. М., Мир, 1976.

[13] Юрьев Д., Комплексная проективная геометрия и квантовая проективная теория поля // ТМФ. 1994. Т.101. вып.3. С.331-348.

[14] Березин Ф.А., Квантование в комплексных симметрических пространствах // Известия АН СССР. Сер.матем. 1975. Т.39, вып.2. С.363-402.

[15] Klimek S., Lesniewski A., Quantum Riemann surfaces. I // Commun.Math.Phys. 1992. V.146. P.103-122.

[16] Биденхарн Л., Лаук Дж., Угловоё момент в квантовоё физике. М., Мир, 1984.

[17] Фукс Д.Б., Когомологии бесконечномерных алгебр Ли. М., Наука, 1984.

[18] Juriev D., Noncommutative geometry, chiral anomaly in the quantum projective ($\mathfrak{sl}(2,\mathbb{C})$–invariant) field theory and $\mathfrak{jl}(2,\mathbb{C})$–invariance // J.Math.Phys. 1992. V.33. P.2819-2822, (E): 1993. V.34. P.1615.

[19] Juriev D., Remarks on nonassociative structures in quantum projective field theory: the

central extension $\mathfrak{jl}(2,\mathbb{C})$ of the double $\mathfrak{sl}(2,\mathbb{C})+\mathfrak{sl}(2,\mathbb{C})$ of the simple Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ and related topics. Acta Appl.Math. 1998. V.50. P.191-196.

[20] Юрьев Д.В., Квантовая проективная теория поля: квантово-полевые аналоги уравнениё Эёлера-Арнольда в проективных G–гипермультиплетах // ТМФ. 1994. Т.98, вып.2. С.220-240.

[21] Мальцев А.И., О представлении неассоциативных колец // УМН. 1952. Т.7, вып.1. С.181-185 [переиздано в “Избранных трудах. 1. Классическая алгебра”, М., Наука, 1976, С.328-331].

[22] Кобаяси Ш., Номидзу К., Основы дифференциальноё геометрии. М., Наука, 1981.

[23] Juriev D., Infinite dimensional geometry and quantum field theory of strings. II. Infinite-dimensional noncommutative geometry of a self-interacting string field // Russian J.Math. Phys. 1996. V.4, no.3. P.287-314.

[24] Juriev D., On the infinite dimensional hidden symmetries. I. Infinite dimensional geometry of q_R–conformal symmetries. E-print: funct-an/9612004 (1996).

[25] Juriev D., On the infinite dimensional hidden symmetries. II. q_R–conformal modular functor. E-print: funct-an/9701009 (1997).

[26] Кириллов А.А., Введение в теорию представлениё и некоммутативныё гармоническиё анализ / Совр.пробл.мат. Фунд.направления. Т.22. М., ВИНИТИ, 1987, С.5-162.

[27] Dubnov V.L., Maslov V.P., Nazaĭkinskiĭ V.E., The complex Lagrangian germ and the canonical operator // Russian J.Math.Phys. 1995. V.3. P.141-180.

[28] Юрьев Д.В., Дроэмы: экспериментальная математика, информатика и бесконечномерная геометрия: Report RCMPI-96/05$^+$ (1996).